

Title	Potential for microbial H ₂ and metal transformations associated with novel bacteria and archaea in deep terrestrial subsurface sediments
Author(s)	Hernsdorf A. W., Amano Yuki, Miyakawa Kazuya, Ise Kotaro, Suzuki Yohei, Anantharaman K., Probst A. J., Burstein David, Thomas B. C., Banfield J. F.
Citation	ISME Journal, 11, p.1915-1929
Text Version	Publisher's Version
URL	https://jopss.jaea.go.jp/search/servlet/search?5055353
DOI	https://doi.org/10.1038/ismej.2017.39
Right	<p>© The Author(s) 2017</p> <p>This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit</p> <p>http://creativecommons.org/licenses/by/4.0/</p>

ORIGINAL ARTICLE

Potential for microbial H₂ and metal transformations associated with novel bacteria and archaea in deep terrestrial subsurface sediments

Alex W Hernsdorf¹, Yuki Amano^{2,3}, Kazuya Miyakawa³, Kotaro Ise², Yohey Suzuki⁴, Karthik Anantharaman⁵, Alexander Probst⁵, David Burstein⁵, Brian C Thomas⁵ and Jillian F Banfield^{5,6}

¹Department of Plant and Microbial Biology, University of California, Berkeley, CA, USA; ²Nuclear Fuel Cycle Engineering Laboratories, Japan Atomic Energy Agency, Tokai, Ibaraki, Japan; ³Horonobe Underground Research Center, Japan Atomic Energy Agency, Horonobe, Hokkaido, Japan; ⁴Graduate School of Science, The University of Tokyo, Tokyo, Japan; ⁵Department of Earth and Planetary Sciences, Berkeley, CA, USA and ⁶Department of Environmental Science, Policy, and Management, Berkeley, CA, USA

Geological sequestration in deep underground repositories is the prevailing proposed route for radioactive waste disposal. After the disposal of radioactive waste in the subsurface, H₂ may be produced by corrosion of steel and, ultimately, radionuclides will be exposed to the surrounding environment. To evaluate the potential for microbial activities to impact disposal systems, we explored the microbial community structure and metabolic functions of a sediment-hosted ecosystem at the Horonobe Underground Research Laboratory, Hokkaido, Japan. Overall, we found that the ecosystem hosted organisms from diverse lineages, including many from the phyla that lack isolated representatives. The majority of organisms can metabolize H₂, often via oxidative [NiFe] hydrogenases or electron-bifurcating [FeFe] hydrogenases that enable ferredoxin-based pathways, including the ion motive Rnf complex. Many organisms implicated in H₂ metabolism are also predicted to catalyze carbon, nitrogen, iron and sulfur transformations. Notably, iron-based metabolism is predicted in a novel lineage of Actinobacteria and in a putative methane-oxidizing ANME-2d archaeon. We infer an ecological model that links microorganisms to sediment-derived resources and predict potential impacts of microbial activity on H₂ consumption and retardation of radionuclide migration.

The ISME Journal (2017) 11, 1915–1929; doi:10.1038/ismej.2017.39; published online 28 March 2017

Introduction

Nuclear energy is deemed to have the potential to offset greenhouse gas emissions from oil and gas power plants and combat climate change, but the wastes generated by this technology may remain highly radioactive for over 100 000 years. Currently, disposal of long-lived radioactive wastes in deep, isolated geological repositories is the only credible long-term solution for human safety (Alexander and McKinley, 2011; Cherry *et al.*, 2014). However, uncertainties exist around the impacts of these repositories on host rock and groundwater, especially over the geologic time frames required for containment. Conversely, the combination of site-specific geology, hydrology, geochemistry and

microbiology are all predicted to provide containment after the eventual breakdown of engineered barrier systems (Harrison *et al.*, 2011; Suzuki *et al.*, 2016).

Microorganisms, ubiquitous components of deep terrestrial ecosystems, thrive on resources such as buried organic carbon, mineral-associated electron donors and acceptors, and biologically or abiotically formed hydrogen and methane (Edwards *et al.*, 2012; Pedersen, 2013). In turn, microbial communities can influence the mineralogical and geochemical features of their environments (Lovley and Chapelle, 1995; Pedersen, 1997; Edwards *et al.*, 2012). Importantly, microbial activities can also impact radioactive waste disposal in the subsurface. For example, microbial respiration can form and maintain reducing conditions following the closure of the disposal facility (Puigdomenech *et al.*, 2001; Amano, *et al.*, 2012a). Additionally, H₂ produced by corrosion of metals associated with repository materials may stimulate H₂-oxidizing bacteria and archaea whose activities may reduce the risks associated with H₂

Correspondence: JF Banfield, Department of Earth and Planetary Sciences, and Department of Environmental Science, Policy, and Management, 369 McCone Hall, Berkeley, CA 94720, USA.
E-mail: jbanfield@berkeley.edu

Received 13 May 2016; revised 2 January 2017; accepted 5 February 2017; published online 28 March 2017

accumulation (Libert *et al.*, 2011; Vinsot *et al.*, 2014; Bagnoud *et al.*, 2016). Some likely H_2 oxidizers include sulfate reducers that generate sulfide that could induce rapid corrosion of containment materials (Rajala *et al.*, 2015), and metal reducers that may reductively immobilize radionuclides (Mohapatra *et al.*, 2010; Williams *et al.*, 2011; Williamson *et al.*, 2013). Furthermore, biofilm development within the pore spaces or fractures of host rocks could affect radionuclide transport and gas diffusion (Pedersen, 2005).

These microbially catalyzed processes also depend on geologic features. For instance, fracture networks in granitic rocks could allow rapid diffusion of gaseous H_2 away from a repository, whereas the low permeability of sedimentary rocks could impede H_2 diffusion (Cherry *et al.*, 2014). Research at the granite-hosted Äspö hardrock laboratory has revealed active microbial H_2 oxidation and sulfate reduction *in situ* as well as uranium immobilization by biofilms (Krawczyk-Bärsch *et al.*, 2012; Pedersen, 2013). Wu *et al.* (2015) recently applied genome-resolved metagenomics to this site to develop a metabolic model of the microbial community, and suggested a higher dependency on organic carbon than previously considered for hardrock environments. Less is known about how microbial communities in sedimentary rock formations may affect repository safety, although research on the Opalinus Clay and Boom Clay formations suggests that the members and complexity of sediment-hosted communities vary immensely by site (Wouters *et al.*, 2013; Bagnoud *et al.*, 2015, 2016).

Understanding microbially mediated processes in sedimentary rocks holds broad significance for global biogeochemical cycling. Deep terrestrial sediments are a vast reservoir of buried organic carbon that is subject to microbial transformation (Breuker *et al.*, 2011; McMahon and Parnell, 2014), often resulting in the formation of carbon dioxide or methane (Pedersen, 2000). Underground research laboratories (URLs) such as Horonobe URL provide the opportunity to study deep subsurface systems. The Horonobe URL has been constructed by the Japan Atomic Energy Agency to conduct basic geoscientific research and evaluate the feasibility and safety of geological disposal in deep sedimentary environments. The URL is situated above Neogene rocks consisting of marine sandstones, mudstones and shales (Hama *et al.*, 2007). Previous microbiological studies at Horonobe URL found a diversity of Proteobacteria and methanogens inhabiting the groundwater, as determined by sequencing of 16S ribosomal RNA (rRNA) gene libraries (Shimizu *et al.*, 2006; Kato *et al.*, 2009). Laboratory studies using rock samples collected from 200 m below ground surface and inoculated with *Pseudomonas denitrificans* supported the development of biofilm communities at *in situ* groundwater conditions (Harrison *et al.*, 2011; Sakurai and Yoshikawa, 2012). However, comprehensive information about the microbial

community structure and metabolic potential of Horonobe groundwaters has not been obtained.

Here, we conducted deep metagenomic sequencing of nine groundwater samples collected from the URL at depths of 140–250 m below the surface. We reconstructed genomes from the sequence data and explored the nature of the subsurface biosphere by incorporating community metabolic potential into a conceptual ecological model. Finally, we used this information to predict how the deep subsurface sediment-associated community could impact radioactive waste repository function in the long term.

Materials and methods

Site description

Groundwater and headspace gas samples were collected from five depths 140–250 m below ground surface at the Horonobe URL (45°02'43"N, 141°51'34"E; Figure 1). At the sampled depths, the rock was composed primarily of soft diatomaceous mudstones, including opal-A and trace amounts of quartz, feldspar, clay minerals, pyrite, calcite and siderite (Hiraga and Ishii, 2007; Ishii *et al.*, 2007; Tachi *et al.*, 2011). The groundwater is 14–18 °C in temperature, brackish and of circumneutral pH. The porewater geochemistry of this site has been determined previously (Sasamoto *et al.*, 2011, 2014; Amano *et al.*, 2012b), and is summarized in Supplementary Table S1. Within the Horonobe URL, three main shafts provide access to subsurface galleries, where boreholes have been drilled into the rock faces. Boreholes sampled during this investigation include 07V140M03 and 09V250M02 drilled from the Ventilation shaft, and three depths within 08E140C01 drilled from the East shaft. All groundwater samples were obtained using a multipacker system developed at the Horonobe URL (Nanjo *et al.*, 2012). To eliminate the influence of drilling and the installation of tools for hydrochemical monitoring (for example, sampling tubes, and casing pipes), groundwaters were drained at least five times the section volume, before sampling for geochemical and microbial studies. The groundwater chemistry has been monitored since 2008, starting immediately after the drilling of these boreholes. Based on the results of geochemical analyses, the compositions of the groundwaters are very stable (Sasamoto *et al.*, 2011, 2014; Amano *et al.*, 2012b). The quality of each groundwater sample was checked by measurement of the concentration of sodium naphthionate, which is an indication of drilling fluid contamination. The concentrations were below the detection limit in all samples.

Headspace gas sampling

Groundwater samples were collected between June 2012 and June 2013, and analyzed for gas composition using a water displacement procedure under




Figure 1 Sampling locations within the Horonobe URL. Abbreviations used in the text follow the format [V/E][Depth][Year], where V/E refers to Ventilation/East Shafts.

atmospheric pressure. The composition of the gas samples (H_2 , N_2 , O_2 , CH_4 , CO_2 and C_2H_6) was determined by gas chromatography. H_2 , N_2 , O_2 and CH_4 were analyzed using a micro-gas chromatograph (GC) (Micro-GC CP-2002; Varian Chrompack International, Agilent Technologies, Middelburg, The Netherlands) packed with a Molecular Sieve-5A and micro-thermal conductivity detector. Oxygen was used as the carrier gas for H_2 analysis, and Ar was used for N_2 , O_2 and CH_4 analyses, as described previously (Miyakawa *et al.*, 2010). CO_2 and C_2H_6 were analyzed using a micro-GC (490 Micro-GC; Agilent Technologies) equipped with a Pora-PLLOTQ column with He as the carrier gas and micro-thermal conductivity detector. The column temperature and pressure of the 490 Micro-GC were maintained at 80 °C and 190 kPa, respectively. Analytical precision was determined by a 95.4% (two sigma) confidence interval of standard gas or air analyses ($n=60$); H_2 is 1.8%, O_2 is 7.9%, N_2 is 9.0%, CH_4 is 8.3%, CO_2 is 2.4% and C_2H_6 is 2.8%.

Biological sampling, DNA extraction and sequencing

Two groundwater samples were collected for metagenomics during 2014 from Ventilation shaft boreholes, one each from boreholes 07V140M03 and 09V250M02. An additional seven samples were collected for metagenomics at two times during 2013 and 2014 from borehole 08E140C01. All samples were collected by passage of groundwater through 0.22 μm membrane filters (type GVWP; Merck Millipore, Darmstadt, Germany). One sample (E215m-2014_f, obtained during 2014 from 215 m depth in borehole 08E140C01) was collected by passage of the 0.22 μm filtrate through a 10 000 nominal molecular weight limit filter (type PLGC; Merck Millipore). Other samples were also collected by this method, but insufficient DNA was recovered for metagenomic analyses. The volume of groundwater samples used for filtration was between 0.9 and 38 L, depending on the cell densities in each groundwater sample. DNA was extracted from the biomass retained on the filters using the Extrap Soil DNA Kit Plus ver. 2 (Nippon Steel and Sumikin Eco-

Tech Corporation, Tsukuba, Japan). Genomic DNA libraries were generated using TruSeq Nano DNA Sample Prep Kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions. Library quality was examined using an Agilent 2100 bioanalyzer (Agilent Technologies) and paired-end 150-bp reads with a 550 bp insert size were sequenced by Hokkaido System Science Co., Ltd (Hokkaido, Japan), using an Illumina HiSeq2000 (San Diego, CA, USA). Sample volumes, DNA concentrations and sequencing information are presented in Supplementary Table S2. All sequencing data were deposited in the NCBI database (BioProject ID: PRJNA321556).

De novo genomic assembly, read mapping and annotation

Raw shotgun sequencing reads were trimmed using the adaptive read trimmer, Sickle (Joshi and Fass, 2011), with default settings. All samples were assembled *de novo* using IDBA-UD with the following parameters: `-mink 40`, `-maxk 100`, `-step 20` and `-pre_correction` (Peng *et al.*, 2012). Trimmed shotgun sequencing reads from each sample were mapped to all scaffolds > 1000 bp, using Bowtie2 with default parameters (Langmead *et al.*, 2009). Further details of the curation process are described in the Supplementary Materials. For all scaffolds over 1000 bp, open reading frames were predicted with Prodigal using the meta setting (Hyatt *et al.*, 2012). Functional annotations for all open reading frames were predicted by BLAST (Altschul *et al.*, 1990) searches against the Uniref100 (Suzek *et al.*, 2007), Uniprot (Magrane and UniProt Consortium, 2011) and KEGG (Kanehisa *et al.*, 2012) databases as described previously (Raveh-Sadka *et al.* 2015). tRNA sequences were predicted using tRNAscan-SE (Schattner *et al.*, 2005), and 16S rRNA sequences were identified using the cmsearch program from the Infernal package (Nawrocki *et al.*, 2009).

Binning and conserved gene analysis

Assembled scaffolds > 1000 bp were binned by a combination of phylogenetic profiles, read coverage and nucleotide content (that is, GC proportion and tetranucleotide signatures) within the binning interface of ggKbase (<http://ggkbase.berkeley.edu/>) and by using emergent self-organizing maps constructed separately from tetranucleotide frequencies and coverage of the scaffolds across the nine different assemblies (Dick *et al.*, 2009; Sharon *et al.*, 2013). Details of the binning procedure are provided in the Supplementary Materials. Resulting bins were evaluated for accuracy and completeness in ggKbase using a set of conserved single-copy phylogenetic marker genes as described previously (Raveh-Sadka *et al.*, 2015; Probst *et al.*, 2016). De-replication of bins was determined using pairwise identity comparisons with Nucmer (Kurtz *et al.*, 2004) via scripts described previously (Probst *et al.*, 2016).

Sequence alignment and phylogeny

Alignment of 16S rRNA genes with reference sequences from the Silva SSU database v.123 was performed with SSU-ALIGN (Nawrocki, 2009) using default parameters, followed by refinement and removal of insertion sequences as described previously (Brown *et al.*, 2015). A set of 16 ribosomal proteins used for phylogenetic placement were identified by profile hidden Markov model (HMM) searches seeded with HMM profiles from the PhyloSift conserved phylogenetic marker genes (Darling *et al.*, 2014). Amino-acid sequences for ribosomal protein were added to a database of ribosomal protein sequences clustered at the genus level (Hug *et al.*, 2016) and aligned using MAFFT (Kato *et al.*, 2002). The alignments were then manually trimmed in Geneious v.8 (BioMatters Ltd., San Francisco, CA, USA) to remove poorly aligned positions at the N and C termini and columns composed of over 90% gaps. The alignments were then concatenated, and sequences with fewer than 50% of all aligned positions were removed before tree construction. Trees were built using RAxML v.8.1.24 (Stamatakis, 2006) on the CIPRES web server (Miller *et al.*, 2010), with the PROTGAMMALG model of evolution and number of bootstraps replicates determined automatically by the extended majority rule.

Taxonomic assignment of bins

All bins were identified to appropriate taxonomic levels in the following order: (i) identification of 16S rRNA genes against the Silva SSU database v.123 (Pruesse *et al.*, 2007); (ii) grouping of concatenated ribosomal protein sequences against the aforementioned database (Hug *et al.*, 2016); (iii) taxonomic clustering of annotated genes by protein BLAST searches against Uniref100, Uniprot and KEGG databases. In cases where taxonomic assignments disagreed among the three methods, bins were assigned to the most specific taxonomic level for which the three methods agreed. Bins in which a majority of the genes had no taxonomic affiliation or that matched viruses were annotated as potential extrachromosomal elements (plasmids and phages).

Estimated coverage of community

We identified a set of unique marker genes by clustering the ribosomal protein S3 (RpS3) assembled across the sampling points. RpS3 was used in place of 16S rRNA genes because of the higher reliability of assembling this gene in metagenomic data sets (Castelle *et al.*, 2013), and its utility as a conserved phylogenetic marker.

Functional gene analysis

Metabolic coding potential of individual bins was explored by reciprocal protein BLAST searches against the aforementioned protein databases and

HMM searches against protein families downloaded from FunGene (Fish *et al.*, 2013), TIGRFAM v.14.0 (Haft *et al.*, 2003) and Pfam v.28.0 (Finn *et al.*, 2014), as well as custom-built profile HMMs for several target genes (Eddy, 2011). Conserved marker genes were used to indicate the potential for autotrophy, carbon degradation and biogeochemical cycles as listed in Supplementary Table S3.

Hydrogenase cofactors were identified as [NiFe], [FeFe] or [Fe] using HMM profiles and aligned with reference sequences from (Greening *et al.*, 2015) for further classification. Group A3 [FeFe] hydrogenases were differentiated from Group A1 by searching for downstream genes required for Group A3 function: *hydB*—a diaphorase ortholog containing multiple [4Fe-4S] clusters; *hydC*—a small protein containing a [2Fe-2S] cluster (Buckel and Thauer, 2013). In contrast, Group A1 [FeFe] hydrogenases are monomeric or have non-catalytic subunits lacking the redox active [4Fe-4S] and [2Fe-2S] clusters found in Group A3 (Greening *et al.*, 2015).

Multiheme cytochromes (MHCs) are predicted to be involved in iron transformations, and can be identified by the conserved heme-binding motif, CXXCH. To estimate the validity of using MHCs as a signal for iron reduction, we downloaded a representative proteome of each prokaryotic genera in the RefSeq database as of 30 October 2015 (Tatusova *et al.*, 2014). Each protein in the reference proteome (and metagenome) was scanned for CXXCH motifs, and labeled as an MHC if it contained 10 or more motifs. Reference genomes containing 4 or more MHCs were investigated to determine if they were from organisms previously identified as capable of dissimilatory iron reduction (Supplementary Table S4).

Results and discussion

Metagenomic reconstruction of groundwater communities

Groundwater samples for metagenomics were obtained from three boreholes at the URL from a depth range of between 140 and 250 m in two consecutive years. Illumina sequencing of the extracted DNA generated a total of 123.6 Gbp of raw paired-end sequence (Supplementary Table S2). We filtered and *de novo* assembled the reads and then binned the assembled sequence using a combination of methods (GC, coverage, taxonomic affiliation, cross-sample abundance and tetranucleotide frequency) to obtain a total of 550 genome bins. De-replication of these bins based on pairwise identity resulted in 228 distinct bins, of which 161 were considered to be high-quality draft genomes (>70% complete; <10% of potential contamination; Supplementary Table S5). The genomes retrieved represent a vast majority of the abundant taxa in the groundwater. Specifically, these genomes represent 49 of the 50 most abundant organisms identified in the East Shaft samples and 46 of the 50 most

abundant organisms identified in the Ventilation Shaft samples (Figure 2). Of the organisms missed by our metagenomic approach, 82% are predicted to comprise <0.1% of the total community, based on sequence abundance information.

Taxonomic assignment of the genomes determined that 15 are from Archaea, 195 are from Bacteria and 19 are from phage or other mobile elements (Figure 3 and Supplementary Figures S1–S3). The communities are highly diverse, with genomic representation of 29 phyla, 13 of which currently lack cultivated representatives and were missed in prior 16S rRNA gene surveys of this site (Shimizu *et al.*,

2006; Kato *et al.*, 2009). Even the genomes from organisms affiliated with relatively well-studied phyla were phylogenetically novel, as over 120 genome bins could not be taxonomically placed beyond the family level (Supplementary Table S5).

Extensive potential for hydrogen metabolism

A notable feature of the genomes was extensive potential for H₂ metabolism, supporting prior indications that H₂ is an important energy currency in the subsurface (Brazelton *et al.*, 2012). A total of 394 hydrogenases were detected in 119 of 161 genomes,




Figure 2 Bars represent the rank abundance curve of 50 most abundant unique *rps3* genes detected in Horonobe assemblies. Red bars indicate scaffolds that were confidently assigned to a genome bin, and black bars indicate scaffolds that remained unbinned. The heights of bars indicate scaffold coverage as a fraction of East shaft (left) and Ventilation shaft (right) communities. Heatmaps show per-sample genome abundance of 10 most abundant genomes from East shaft (left) and Ventilation shaft (right).




Figure 3 Phylogenetic tree of 130 high-quality genomes constructed from concatenated sequences of 16 ribosomal proteins. Sequences were excluded from the tree if they contained fewer than 450 aligned positions. Support for internal nodes was constructed from 100 bootstrap replicates (white ≥ 50%, gray ≥ 75%, black ≥ 95% confidence, no shading ≤ 50%). Stars represent essentially complete genomes.

and across 20 phyla (Figure 4). To evaluate the functional role and directionality of these genes in H_2 metabolism, we identified the cofactor (that is, [NiFe], [FeFe] or [Fe]) and classified each hydrogenase using HMM profiles and phylogenetic association with reference databases (Greening *et al.*, 2015).

[FeFe] hydrogenases were distinguished into three major groups (A, B and C) by sequence identity and Group A was further subdivided based on operon structure, as phylogeny alone was insufficient to distinguish the function of [FeFe] Group A hydrogenases. Surprisingly, ~80% of the [FeFe] Group A hydrogenases were identified as heterotrimeric/heterotetrameric Group A3 [FeFe] hydrogenases, which reversibly bifurcate electrons from H_2 to ferredoxin and NAD (Greening *et al.*, 2015). Another 15% were of monomeric composition (Group A1), whereas the remaining 5% could not be reliably assigned to A1 or A3 because the genes were located adjacent to the end of the scaffold. The high occurrence of bifurcating hydrogenases found at Horonobe has not been recognized in studies at other sites, although bioinformatics studies have suggested that a significant fraction of H_2 -producing anaerobes harbor bifurcating hydrogenases (Schut and Adams, 2009; Greening *et al.*, 2015). However, a comparison of the abundance of bifurcating and monomeric [FeFe] hydrogenases in environmental systems has not been possible until recently, as genomic context is

required to differentiate the function of the catalytic subunits.

Although CH_4 and CO_2 comprised a vast majority (>99%) of gases obtained from all the sampling zones (Table 1), H_2 levels were also consistently elevated across the sampling zones (11.6–36.4 p.p.m.) during a multiyear sampling effort. The high levels of H_2 in the Horonobe subsurface make H_2 oxidation a favorable strategy for generating reduced cofactors like ferredoxin and NAD, and has likely enriched for organisms with genomes containing bifurcating hydrogenases. In support of this, Rnf complexes were detected in over 70% of the genomes with bifurcating [FeFe] hydrogenases, which could allow the direct coupling of hydrogen oxidation to generation of a proton gradient for ATP synthase (Biegel *et al.*, 2011; Tremblay *et al.*, 2013). Additionally, the few monomeric, H_2 -evolving [FeFe] hydrogenases (Group A1) that were identified nearly always co-occurred with bifurcating hydrogenases (Figure 5 and Supplementary Figure S4). This high degree of co-occurrence suggests that organisms that evolve H_2 (that is, those that have A1 hydrogenases) may also rely upon uptake hydrogenases (that is, A3 hydrogenases) to fuel their metabolism in times of substrate limitation. In previous metagenomic surveys of H_2 -metabolizing communities, [FeFe] hydrogenases have been frequently taken as an indication of H_2 production (Brazelton *et al.*, 2012). Indeed, the *in situ* direction of bifurcating hydrogenases cannot be




Figure 4 Number of hydrogenases detected in the Horonobe genomes, by phylum and class, in the case of Proteobacteria. Hydrogenase type is annotated as [Cofactor]-Group-Localization (M = membrane-bound, P = periplasmic, C = cytoplasmic)-Function.

Table 1 Headspace gas composition of groundwater samples

	V140m, n = 10	E160m, n = 14	E185m, n = 11	E215m, n = 12	V250m, n = 13
CH_4 (%)	88.8 (± 3.4)	85.5 (± 8.0)	97.7 (± 3.4)	99.4 (± 1.8)	88.6 (± 1.7)
CO_2 (%)	10.5 (± 3.0)	13.8 (± 6.3)	2.8 (± 2.6)	1.2 (± 0.5)	9.6 (± 3.3)
N_2 (%)	0.50 (± 0.54)	0.49 (± 0.30)	0.46 (± 0.16)	0.44 (± 0.10)	0.58 (± 0.34)
O_2 (%)	0.04 (± 0.03)	0.06 (± 0.06)	0.05 (± 0.07)	0.03 (± 0.02)	0.05 (± 0.04)
H_2 (p.p.m.)	11.6 (± 22.5)	33.0 (± 58.7)	36.4 (± 90.2)	32.8 (± 46.1)	22.7 (± 38.6)
C_2H_6 (p.p.m.)	54 (± 29)	40 (± 30)	39 (± 30)	31 (± 29)	44 (± 36)

Values presented indicate the mean and standard deviation of measurements taken approximately once a month between June 2012 and June 2013.




Figure 5 Metabolic potential to influence biogeochemical cycles of 59 genomes with an average abundance >0.1% across all samples. Differentially shaded tiles in the %Ave column represent average relative abundance of the genomes as a fraction of the complete community. Differentially shaded tiles for fermentation, complex carbon degradation and hydrogenases indicate the relative number of genes involved in such processes.

determined from sequence alone, as they could function in the reverse direction as confurcating hydrogenases (Sieber *et al.*, 2012). However, the abundance of bifurcating [FeFe] hydrogenases and co-occurrence of Group A1 with A3 hydrogenases suggests that the presence of [FeFe] hydrogenases alone does not indicate H₂ production. Similar patterns have also been observed recently in studies of human stool metagenomes (Wolf *et al.*, 2016). These results have important implications when interpreting the results of read-based, rather than genome-based metagenomics.

[FeFe] Group B hydrogenases, whose function is currently unknown, were also identified in 17 genomes, but were found exclusively in organisms that also harbored bifurcating Group A3 hydrogenases. Group C putative sensory [FeFe] hydrogenases were likewise abundant and only identified in genome bins that also had genes for H₂ uptake or evolution. The high incidence of Group B and C [FeFe] hydrogenases from genomes at Horonobe suggest these enzymes are highly important in environmental systems, and points to the need to further characterize this prevalent group.

A total of 155 [NiFe] hydrogenases were identified across the four major [NiFe] hydrogenase groups, most of which were found in groups commonly considered to have uptake or bidirectional functions. As with the [FeFe] hydrogenases, the genomic evidence for H₂ uptake in [NiFe] hydrogenases reflects the value of H₂ as a source of reducing power in the deep subsurface and suggests that elevated H₂ concentrations have enriched for hydrogenotrophs.

Metals cycling catalyzed by novel lineages

Sediment-associated ferric iron-bearing minerals, including smectite and chlorite, and mineral coatings also provide potential resources for microbial metabolism. To evaluate the microbial potential to reduce such minerals, we searched the genomes for MHCs (operationally defined as cytochromes with 10 or more heme-binding motifs). MHCs have been identified as components of electron transferring complexes in metal-reducing bacteria like *Geobacter sulfurreducens* and *Shewanella oneidensis* (Weber *et al.*, 2006), and may be general predictors of iron reduction capacity given that organisms with many MHCs are almost exclusively iron-reducing bacteria and archaea. By scanning the RefSeq genome database for MHCs, we found that 1.2% of the unique species in the database contained 4 or more MHCs, of which 90% have been shown or predicted to be capable of dissimilatory iron reduction (Supplementary Table S5). Using these criteria on our metagenome dataset, we identified 253 MHCs from 63 genomes, and predict that 23 genomes encode the capacity for iron reduction (Figure 5 and Supplementary Figure S4).

Interestingly, the genome of the most abundant organism in the E215m zone is an ANME-2d archaeon (HGW-Methanoperedenaceae-1) that encodes 11 MHCs, four of which include 50 or more heme-binding motifs. To our knowledge, these represent the most heme-rich cytochromes sequenced to date in any organism. Although other ANME lineages have recently been hypothesized to use MHCs for direct transfer of electrons to syntrophic Deltaproteobacteria partners (McGlynn *et al.*, 2015), the abundance of HGW-Methanoperedenaceae-1 across our samples is not significantly correlated with any other organism (using an abundance threshold of 10 : 1). The HGW-Methanoperedenaceae-1 genome does not encode the ability to use nitrate or sulfate as electron acceptors, although it has a complete methanogenesis/methane-oxidation pathway. Thus, we suggest that the MHCs are used in the direct reduction of iron minerals coupled to methane oxidation. These predictions are further supported by multivariate functional analyses that validated the correlation between total Fe in groundwater and the abundance of HGW-Methanoperedenaceae-1 (Supplementary Figure S5). The inference is consistent with a very recent report linking ANME-2d archaea to iron reduction in a freshwater enrichment culture from a Dutch canal (Ettwig *et al.* 2016).

Other organisms that were enriched in the E215m zone at lower abundance included several H₂ oxidizing iron reducers. Dissimilatory metal reduction coupled to H₂ oxidation is often associated with Gram-negative bacteria (for example, species of *Geobacter* and *Shewanella*), and indeed, a majority of the genomes capable of metal reduction in the RefSeq database are Proteobacteria. However, Firmicutes of the Peptococcaceae family also use this metabolism (Junier *et al.*, 2010; Wrighton *et al.*, 2011). Interestingly, the four Horonobe genomes containing the most MHCs belong to Gram-positive organisms whose genomes also harbor one or more membrane-bound H₂ uptake hydrogenases [NiFe], which can liberate electrons for metals respiration (Greening *et al.*, 2015). The genome with the most MHCs is from a member of the Peptococcaceae, but surprisingly, the other three genomes were all from organisms distantly related to isolated members of the Actinobacteria. Currently isolated Actinobacteria capable of dissimilatory iron reduction are obligate acidophiles and lack MHCs in their genomes (Bridge and Johnson, 1998; Itoh *et al.*, 2011; Stackebrandt, 2014). The Horonobe clade diverges deeply from Coriobacteriobacteriales isolates, although their 16S rRNA gene sequences are closely related to those detected in iron enrichment cultures sampled from pond sediments and rice paddy soils (Supplementary Figure S6; Wang *et al.*, 2009; Lentini *et al.*, 2012). These genomes may represent an important group of neutrophilic iron-reducing organisms that has previously gone unnoticed.

Sulfur cycling

We explored the potential for sulfur cycling by looking for key genes in pathways for sulfate reduction and oxidation of reduced sulfur compounds. In total, 20 organisms, generally representatives from the Betaproteobacteria, are predicted to conserve energy from the oxidation of reduced sulfur species, including sulfide, elemental sulfur and thiosulfate (Figure 5 and Supplementary Figure S4). An additional 15 organisms are capable of reducing sulfate; all are affiliated with the Deltaproteobacteria, Firmicutes and Rifebacteria (previously known as ACD39; Anantharaman *et al.*, 2016).

The abundance of organisms capable of sulfur cycling was not anticipated from the groundwater geochemistry, given that sulfate and sulfide concentrations in Horonobe groundwaters are consistently below 3 μM . However, analysis of porewaters extracted directly from samples of the rock matrix revealed levels of sulfate that were at least 2 orders of magnitude higher than in the groundwater (Kunimaru *et al.*, 2010). Although the high levels of sulfate detected in the porewaters were attributed to sample disturbance via oxidation of pyrite (Sasamoto *et al.*, 2011), microbially catalyzed oxidation of pyrite could liberate free sulfate, which is likely to be rapidly scavenged. The abundance of organisms capable of oxidizing and reducing sulfur species may support a cryptic sulfur cycle, where the active cycling of sulfur between solid and dissolved forms keeps concentrations low.

Nitrogen cycling

Evidence for nitrogen cycling was investigated by searching the genomes for genes involved in nitrogen fixation, nitrate reduction, nitrification and anammox. Nitrogen fixation potential was detected in 37 genomes, which together comprise roughly one-third of the community by abundance and include four of the ten most abundant genomes from the East shaft (Figure 5 and Supplementary Figure S4).

Nitrate reduction is predicted to be a common metabolic capacity, with 72 genomes encoding at least one step in the nitrate reduction pathway. However, the pathway was frequently found to be incomplete, as only nine genomes have all the genes necessary to complete the denitrification pathway to N_2 , and 19 are capable of dissimilatory nitrite reduction to ammonia. Significant overlap exists between these two pathways, and 13 genomes are capable of reducing nitrite using either pathway.

In contrast to the number of genomes predicted to encode nitrate reduction pathways, only four genomes were predicted to encode partial nitrification pathways (each genome has a hydroxylamine dehydrogenase homolog, but lacks ammonia monooxygenase), suggesting that nitrification is not a major energy-conserving pathway in Horonobe sediments. Furthermore, hydroxylamine dehydrogenase has been shown to catalyze the reverse reaction

in vitro, and so may be acting as a means of detoxification when nitrite levels are elevated (Kostera *et al.*, 2008). No copies of the hydrazine oxidoreductase gene were identified, and so anammox is likewise not predicted to be a capacity of any of the studied organisms.

Given that nitrate reduction capacity is frequently encoded in the studied genomes, nitrate-reducing microorganisms are likely capable of scavenging this compound at low levels. The activity of such microorganisms has resulted in depletion of nitrate, to the extent that nearly all nitrogen in both groundwater and porewater samples was present as N_2 or ammonia. Further, the persistence of ammonia may be the consequence of a microbial biosphere in which nitrification pathways are essentially absent.

Carbon cycling

Genes involved in carbon fixation were identified in roughly one quarter of the genome bins, representing 43 and 77% of the East shaft and Ventilation shaft communities by abundance (Figure 5 and Supplementary Figure S4). Notably, the groundwater samples collected from Ventilation shaft were dominated by an archaeon from the recently described order Altiarchaeales (Probst *et al.*, 2014). The presence of this organism would have been missed in prior 16S rRNA gene surveys of Horonobe groundwaters as commonly used primer pairs fail to amplify the 16S rRNA gene of this group (Rudolph *et al.*, 2001). Altiarchaeales were predicted to use the Wood–Ljungdahl for carbon fixation (Probst *et al.*, 2014). This pathway was the most frequently encoded strategy for carbon fixation, detected in 24 genomes mainly from Deltaproteobacteria, Firmicutes, and Archaea, including the methanogens. The abundance of autotrophs using the Wood–Ljungdahl pathway has also been found in another deep terrestrial environment (Magnabosco *et al.*, 2015), suggesting that this pathway may be an important link in the deep carbon cycle.

The methanogen genomes are all predicted to couple CO_2 reduction to hydrogenotrophy based on the uptake hydrogenases in their genomes, and none were capable of using acetate or methanol for methanogenesis. Each methanogen genome also harbors one or more formate dehydrogenase homologs, and so may be capable of oxidizing formate produced via fermentative reactions, although the capacity for formate utilization cannot be predicted solely by the presence of formate dehydrogenase (Wood *et al.*, 2003). Regardless of the electron donor, methanogenesis is predicted to be the predominant source of methane in these depths, based on carbon and hydrogen isotope ratios of methane measured in dissolved gases (Tamamura *et al.*, 2014). This biogenic methane likely supports the growth of anaerobic methane oxidizers, such as HGW-Methanoperedenaceae-1.

The only other carbon fixation pathway identified was the Calvin cycle, as determined by the presence of RuBisCO. Interestingly, one complete (curated, closed) genome from a member of the recently described Candidate Phyla Radiation (Brown *et al.*, 2015), HGW-Dojkabacteria-1 (the second complete genome of this phylum previously referred to as WS6; Wrighton *et al.*, 2016), was found to harbor genes for both form II/III and form III-like RuBisCO that are implicated in the archaeal-type nucleotide salvage pathway, a phenomenon not reported previously. This result is surprising, given that the genome is <800 kbp in length, and suggests the importance of CO₂ incorporation into nucleotide-derived sugars in the metabolism of this organism.

Heterotrophic lifestyles were inferred based on a lack of genes for CO₂ fixation and presence of genes for organic carbon degradation. Based on a lack of genes for the TCA cycle, electron transport chain or other respiratory enzymes, the most abundant organism detected in the East Shaft, HGW-Saccharibacteria-1 (from the Candidate Phyla Radiation phylum previously referred to as TM7), is predicted to be a fermenter. HGW-Saccharibacteria-1 is also likely dependent on the groundwater community for survival, due to the near-complete lack of essential biosynthetic pathways (for example, for biosynthesis of amino acids, nucleotides and lipids) in its genome, a prediction that is consistent with those for other representatives of this group (Kantor *et al.*, 2013). A similar lifestyle is predicted for a bacterium affiliated with the Peregrinibacteria (a Candidate Phyla Radiation phylum) for which we reconstructed a complete (curated, closed) genome.

Complex carbon (for example, cellulose, chitin, lignin and aromatic amino acids) degradation was a common functional trait predicted for the Horonobe organisms (Supplementary Table S6). Over half the draft genomes encode genes predicted to degrade one or more forms of cellulose, hemicellulose and chitin. Cellulolytic- and chitin-degrading enzymes were particularly common in Chloroflexi, Bacteroidetes and Ignavibacteria, although genomes from several poorly characterized phyla also had extensive capacity to degrade complex carbon. In particular, a genome from a previously unrecognized phylum encodes 12 putative cellulases. We propose that the name 'Goldbacteria' is reserved for this phylum, should further discoveries support the inference that this is a distinct major lineage in Domain Bacteria. This suggestion is in recognition of Thomas Gold's contributions to the study of the deep biosphere. The genome also has multiple cellobiases and other hemicellulose and endohemicellulose-debranching enzymes, making it a likely degrader of plant-derived organics. *N*-alkane monooxygenases were also detected in 12 Betaproteobacteria and Gammaproteobacteria, along with pathways for degradation of aromatic compounds, including nitrobenzene, aniline, benzoate, naphthalene and toluene. Notably, five of these genomes were from

organisms most closely related to the known aromatic compound-degrading organism, *Dechloromonas aromatica* (Coates *et al.*, 2001), and each maintain several pathways for breakdown of benzene via catechol cleavage by catechol 2,3-dioxygenase. However, given the lack of oxygen in the Horonobe subsurface, the organisms related to *D. aromatica* may rely on an alternative mechanism for degrading aromatic compounds, although the genes involved in this pathway have yet to be identified (Salinero *et al.*, 2009).

Genes and pathways involved in degradation of complex carbon compounds likely degrade recalcitrant organic materials trapped during sedimentation (Ishii *et al.*, 2011). In particular, the silica skeletons of diatoms buried in sediments may have locked away organics that provide carbon for microbial metabolism for millions of years (Parkes *et al.*, 2014). Additionally, the salinity profiles in the vicinity of the boreholes suggest dilution of the original seawater with freshwater (Hama *et al.*, 2007), which could bring surface-derived organics, such as humic substances, into contact with deep groundwater communities. Intermittent lack of access to such resources and the availability of dissolved inorganic carbon may also select for autotrophic organisms that continue to add to the carbon pool.

A conceptual ecological model of the deep sedimentary subsurface

Taken together, these genome-based metabolic predictions provide insight into the metabolic foundation for the biosphere in a deep sedimentary environment. According to the previous studies on the mineralogy of diatomaceous siliceous sedimentary rocks in the Horonobe area, the dominant minerals are quartz, feldspars, pyrite and carbonate (siderite and/or magnesite). Also present are clay minerals, including kaolinite, smectite, illite and/or chlorite (Hiraga and Ishii, 2007; Ishii *et al.*, 2007; Tachi *et al.*, 2011). Modeling of groundwater geochemistry using PhreeqC predicts that some Fe-rich clay minerals, such as smectite, nontronite and daphnite, are oversaturated, as are pyrite, Fe(OH)₃ and goethite (Supplementary Table S7). These calculations are consistent with the observed mineralogy of rock core samples from the Horonobe site. The previously reported geological, geochemical and mineralogical information, combined with microbial results obtained here, were integrated into a conceptual model to describe the links between sediment resources and microbial ecology (Figure 6).

The abundant organic carbon buried in diatomaceous sediment provides fuel for respiratory organisms (Figures 6a and e) and fermenters (Figure 6h), which generate CO₂, H₂ and simple organic compounds. The buried organic materials are also likely to be an important source of nitrogen compounds, as N₂ gas concentrations are extremely depleted in




Figure 6 A proposed ecological model of the Horonobe subsurface. Labeled microbial populations are described in the main text.

comparison with the Earth's atmosphere. We infer that biomass-derived nitrogen is assimilated, as there is no evidence for ammonia oxidation capacity in any organism detected from any sampled site. This apparently precludes operation of a complete nitrogen cycle.

H₂ and organics produced by fermentation can serve as electron donors to iron (Figure 6e) and sulfate (Figure 6a) reducers. Sulfate reducers depend on the availability of sulfate, which may be derived from gypsum (Figure 6a) formed, in some cases, by pyrite oxidation occurring during the construction of a repository (Figures 6b and c). However, gypsum was undersaturated in all groundwaters, based on geochemical modeling (Supplementary Table S7), and it is very rarely detected in sediments. Therefore, sulfate reduction is probably limited by low availability of sulfate. H₂ is also capable of fueling carbon fixation (Figures 6b, c and g), which can provide a feedback loop for regeneration of organic carbon. H₂ is also used as a substrate for methanogenesis (Figure 6f). Anaerobic methane oxidation may be coupled to iron reduction, as suggested previously (Beal *et al.*, 2009; Chang *et al.*, 2012; Ettwig *et al.*, 2016). We infer that methane-oxidation by ANME-2d archaea is coupled to reduction of ferric iron derived from sediment-associated iron-bearing minerals.

Conclusions

This study has substantially increased genomic resolution compared with prior deep subsurface studies, with draft genomes for 80% of detected organisms. This level of genomic sampling enabled a detailed analysis of metabolic potential and pathway configurations in the sediment-associated microbial communities. The results uncovered a variety of interesting metabolic capacities, including large MHCs likely indicative of iron reduction in both Actinobacteria and methane-oxidizing archaea. We also found evidence of fermentation-based metabolisms in organisms inferred to be obligate symbionts of bacteria, and no evidence for the oxidative branch of the nitrogen cycle. Hydrogen-based metabolism, often mediated by [FeFe] bifurcating hydrogenases and Rnf

complexes, is widely distributed and likely central to microbial community function.

The results reveal possible feedbacks into long-term safety of geological disposal in sedimentary environments. Elevated H₂ levels may develop during storage of radioactive waste in some geological repositories. This gas may stimulate activity of hydrogenotrophic organisms, reducing the potential for gas accumulation in the low-diffusivity sedimentary rock environment (Vinsot *et al.*, 2014). Increased H₂ is expected to stimulate organisms that interact directly with other biogeochemical cycles. Numerous hydrogenotrophic sulfate reducers are likely to produce sulfide. Even though dissolved concentrations of sulfur and sulfide may be low initially, construction of a geological repository involves a significant disturbance of local groundwater conditions and heat from the radioactive decay of buried wastes, and will likely drive extensive groundwater circulation. These disturbances could result in oxidation of FeS₂ minerals and liberation of sulfate into the groundwater. Alternatively, sulfate could be supplied from distantly located sources, including oxidized zones closer to the surface. The microbial production of sulfide would accelerate the corrosion of radioactive waste containers (Rajala *et al.*, 2015). Hydrogenotrophic iron-reducing organisms may also affect the migration of radionuclides in the sedimentary rock environments (Wall and Krumholz, 2006; Junier *et al.*, 2009). The abundance of MHCs in the genomes supports the potential for enzymatic reduction of U(VI) (Marshall *et al.*, 2006) and abiotic reduction of U(VI) by biologically reduced iron (Wu *et al.*, 2010). ⁹⁹Tc, a long-lived fission product of ²³⁵U, can likewise be reduced by biogenic Fe(II) and enzymatic activity of [NiFe] hydrogenases (De Luca *et al.*, 2001). Laboratory and field investigations have established that the subsurface migration of long-lived radionuclides is profoundly influenced by such enzymatic pathways (Suzuki *et al.*, 2002; Wu *et al.*, 2010; Williams *et al.*, 2011). Stimulation of metal-reducing communities could likewise retard the movement of dissolved radionuclides during the lifetime of the repository. The results presented here elucidate the potential for microbial communities to affect geochemical processes relevant for nuclear

waste disposal in deep sedimentary environments and the role of hydrogen in both potentially positive and negative processes. The findings also provide insight regarding the microbiology of a deep subsurface sedimentary environment and identify potential metabolic roles for a variety of relatively little known bacteria and archaea.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

This material is based on work supported by the National Science Foundation Graduate Research Fellowship under Grant No. (DGE 1106400). This study was performed as a part of 'The project for validating assessment methodology in geological disposal system' funded by the Ministry of Economy, Trade and Industry of Japan.

References

- Alexander WR, McKinley L. (2011). *Deep Geological Disposal of Radioactive Waste*, Vol. 9. Elsevier: Amsterdam.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- Amano Y, Sasao E, Niizato T, Iwatsuki T. (2012a). Redox buffer capacity in water–rock–microbe interaction systems in subsurface environments. *Geomicrobiol J* **29**: 628–639.
- Amano Y, Yamamoto Y, Nanjyo I, Murakami H, Yokota H, Yamazaki M *et al.* (2012b). Data of groundwater from boreholes, river water and precipitation for the Horonobe Underground Research Laboratory Project (2001–2010). Available at: <http://jolissrch-inter.tokai-sc.jaea.go.jp/pdfdata/JAEA-Data-Code-2011-023.pdf>.
- Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ *et al.* (2016). Metabolic handoffs shape biogeochemical cycles mediated by complex microbial communities. *Nat Commun* **7**: 13219.
- Bagnoud A, Chourey K, Hettich RL, de Bruijn I, Andersson AF, Leupin OX *et al.* (2016). Reconstructing a hydrogen-driven microbial metabolic network in Opalinus Clay rock. *Nat Commun* **7**: 12770.
- Bagnoud A, de Bruijn I, Andersson AF, Diomidis N, Leupin OX, Schwyn B *et al.* (2015). A minimalistic microbial food web in an excavated deep subsurface clay rock. *FEMS Microbiol Ecol* **92**: fiv138.
- Beal EJ, House CH, Orphan VJ. (2009). Manganese- and iron-dependent marine methane oxidation. *Science* **325**: 184–187.
- Biegel E, Schmidt S, González JM, Müller V. (2011). Biochemistry, evolution and physiological function of the Rnf complex, a novel ion-motive electron transport complex in prokaryotes. *Cell Mol Life Sci* **68**: 613–634.
- Brazelton WJ, Nelson B, Schrenk MO. (2012). Metagenomic evidence for H₂ oxidation and H₂ production by serpentinite-hosted subsurface microbial communities. *Front Microbiol* **2**: 1–16.
- Breuker A, Köweker G, Blazejak A, Schippers A. (2011). The deep biosphere in terrestrial sediments in the Chesapeake Bay area, Virginia, USA. *Front Microbiol* **2**: 1–13.
- Bridge TAM, Johnson DB. (1998). Reduction of soluble iron and reductive dissolution of ferric iron-containing minerals by moderately thermophilic iron-oxidizing bacteria. *Appl Environ Microbiol* **64**: 2181–2186.
- Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A *et al.* (2015). Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523**: 208–211.
- Buckel W, Thauer RK. (2013). Energy conservation via electron bifurcating ferredoxin reduction and proton/Na⁺ translocating ferredoxin oxidation. *Biochim Biophys Acta* **1827**: 94–113.
- Castelle CJ, Hug LA, Wrighton KC, Thomas BC, Williams KH, Wu D *et al.* (2013). Extraordinary phylogenetic diversity and metabolic versatility in aquifer sediment. *Nat Commun* **4**: 2120.
- Chang Y-H, Cheng T-W, Lai W-J, Tsai W-Y, Sun C-H, Lin L-H *et al.* (2012). Microbial methane cycling in a terrestrial mud volcano in eastern Taiwan. *Environ Microbiol* **14**: 895–908.
- Cherry JA, WiM Alley, Parker BL. (2014). Geologic disposal of spent nuclear fuel: an earth science perspective. *Bridg* **44**: 51–59.
- Coates JD, Chakraborty R, Lack JG, O'Connor SM, Cole KA, Bender KS *et al.* (2001). Anaerobic benzene oxidation coupled to nitrate reduction in pure culture by two strains of *Dechloromonas*. *Nature* **411**: 1039–1043.
- Darling AE, Jospin G, Lowe E, Matsen FAI, Bik HM, Eisen JA. (2014). PhyloSift: phylogenetic analysis of genomes and metagenomes. *Peer J* **2**: e243.
- Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP *et al.* (2009). Community-wide analysis of microbial genome sequence signatures. *Genome Biol* **10**: R85.
- Eddy SR. (2011). Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**: e1002195.
- Edwards KJ, Becker K, Colwell F. (2012). The deep, dark energy biosphere: intraterrestrial life on Earth. *Annu Rev Earth Planet Sci* **40**: 551–568.
- Ettwig KF, Zhu B, Speth D, Keltjens JT, Jetten MSM, Kartal B. (2016). Archaea catalyze iron-dependent anaerobic oxidation of methane. *Proc Natl Acad Sci USA* **113**: 12792–12796.
- Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR *et al.* (2014). Pfam: the protein families database. *Nucleic Acids Res* **42**: D222–D230.
- Fish JA, Chai B, Wang Q, Sun Y, Brown CT, Tiedje JM *et al.* (2013). FunGene: the functional gene pipeline and repository. *Front Microbiol* **4**: 00291.
- Greening C, Biswas A, Carere CR, Jackson CJ, Taylor MC, Stott MB *et al.* (2015). Genomic and metagenomic surveys of hydrogenase diversity indicate H₂ is a widely-utilised energy source for microbial growth and survival. *ISME J* **10**: 761–777.
- Haft DH, Selengut JD, White O. (2003). The TIGRFAMs database of protein families. *Nucleic Acids Res* **31**: 371–373.
- Hama K, Kunimaru T, Metcalfe R, Martin AJ. (2007). The hydrogeochemistry of argillaceous rock formations at the Horonobe URL site, Japan. *Phys Chem Earth* **32**: 170–180.

- Harrison H, Wagner D, Yoshikawa H, West JM, Milodowski AE, Sasaki Y *et al.* (2011). Microbiological influences on fracture surfaces of intact mudstone and the implications for geological disposal of radioactive waste. *Miner Mag* **75**: 2449–2466.
- Hiraga N, Ishii E. (2007). Mineral and chemical composition of rock core and surface gas composition in Horonobe Underground Research Laboratory Project (Phase 1). JAEA-Data/Code 2007-022. Available at: <http://jolissrch-inter.tokai-sc.jaea.go.jp/pdfdata/JAEA-Data-Code-2007-022.pdf>.
- Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ *et al.* (2016). A new view of the tree and life's diversity. *Nat Microbiol* **1**: 16048.
- Hyatt D, LoCascio PF, Hauser LJ, Uberbacher EC. (2012). Gene and translation initiation site prediction in metagenomic sequences. *Bioinformatics* **28**: 2223–2230.
- Ishii E, Hama K, Kunimaru T, Sato H. (2007). Change in groundwater pH by infiltration of meteoric water into shallow part of marine deposits. *J Geol Soc Jpn* **113**: 41anSo.
- Ishii E, Sanada H, Iwatsuki T, Sugita Y, Kurikami H. (2011). Mechanical strength of the transition zone at the boundary between opal-A and opal-CT zones in siliceous rocks. *Eng Geol* **122**: 215–221.
- Itoh T, Yamanoi K, Kudo T, Ohkuma M, Takashina T. (2011). *Aciditerrimonas ferrireducens* gen. nov., sp. nov., an iron-reducing thermoacidophilic actinobacterium isolated from a solfataric field. *Int J Syst Evol Microbiol* **61**: 1281–1285.
- Joshi N, Fass J. (2011). Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. Available at: <https://github.com/najoshi/sickle>.
- Junier P, Fruttschi M, Wigginton NS, Schofield EJ, Bargar JR, Bernier-Latmani R. (2009). Metal reduction by spores of *Desulfotomaculum reducens*. *Environ Microbiol* **11**: 3007–3017.
- Junier P, Junier T, Podell S, Sims DR, Detter JC, Lykidis A *et al.* (2010). The genome of the Gram-positive metal- and sulfate-reducing bacterium *Desulfotomaculum reducens* strain MI-1. *Environ Microbiol* **12**: 2738–2754.
- Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. (2012). KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res* **40**: D109–D114.
- Kantor RS, Wrighton KC, Handley KM, Sharon I, Hug LA, Castelle CJ *et al.* (2013). Small genomes and sparse metabolisms of sediment-associated bacteria from four candidate phyla. *MBio* **4**: 1–11.
- Kato K, Nagaosa K, Kimura H, Katsuyama C, Hama K, Kunimaru T *et al.* (2009). Unique distribution of deep groundwater bacteria constrained by geological setting. *Environ Microbiol Rep* **1**: 569–574.
- Katoh K, Misawa K, Kuma K, Miyata T. (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**: 3059–3066.
- Kostera J, Youngblut M, Slosarczyk J, Pacheco AA. (2008). Kinetic and product distribution analysis of NO^{*} reductase activity in *Nitrosomonas europaea* hydroxylamine oxidoreductase. *J Biol Inorg Chem* **13**: 1073–1083.
- Krawczyk-Bärsch E, Lünsdorf H, Pedersen K, Arnold T, Bok F, Steudtner R *et al.* (2012). Immobilization of uranium in biofilm microorganisms exposed to groundwater seeps over granitic rock tunnel walls in Olkiluoto, Finland. *Geochim Cosmochim Acta* **96**: 94–104.
- Kunimaru T, Ota K, Alexander WR, Yamamoto H. (2010). Groundwater/Porewater Hydrochemistry at Horonobe URL: Data Freeze I. Available at: <http://jolissrch-inter.tokai-sc.jaea.go.jp/pdfdata/JAEA-Research-2010-035.pdf>.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C *et al.* (2004). Versatile and open software for comparing large genomes. *Genome Biol* **5**: R12.
- Langmead B, Trapnell C, Pop M, Salzberg SL. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**: R25.
- Lentini CJ, Wankel SD, Hansel CM. (2012). Enriched iron (III)-reducing bacterial communities are shaped by carbon substrate and iron oxide mineralogy. *Front Microbiol* **3**: 00404.
- Libert M, Bildstein O, Esnault L, Jullien M, Sellier R. (2011). Molecular hydrogen: an abundant energy source for bacterial activity in nuclear waste repositories. *Phys Chem Earth* **36**: 1616–1623.
- Lovley DR, Chapelle FH. (1995). Deep subsurface microbial processes. *Rev Geophys* **33**: 365–381.
- De Luca G, De Philip P, Dermoun Z, Rousset M, Vermeglio A. (2001). Reduction of Technetium(VII) by *Desulfovibrio fructosovorans* is mediated by the nickel-iron hydrogenase. *Appl Environ Microbiol* **67**: 4583–4587.
- Magnabosco C, Ryan K, Lau MCY, Kuloyo O, Sherwood Lollar B, Kieft TL *et al.* (2015). A metagenomic window into carbon metabolism at 3 km depth in Precambrian continental crust. *ISME J* **10**: 730–741.
- Magrane M, UniProt Consortium (2011). UniProt Knowledgebase: a hub of integrated protein data. *Database* **2011**: bar009.
- Marshall MJ, Beliaev AS, Dohnalkova AC, Kennedy DW, Shi L, Wang Z *et al.* (2006). c-Type cytochrome-dependent formation of U(IV) nanoparticles by *Shewanella oneidensis*. *PLoS Biol* **4**: e268.
- McGlynn SE, Chadwick GL, Kempes CP, Orphan VJ. (2015). Single cell activity reveals direct electron transfer in methanotrophic consortia. *Nature* **526**: 531–535.
- Mcmahon S, Parnell J. (2014). Weighing the deep continental biosphere. *FEMS Microbiol Ecol* **87**: 113–120.
- Miller MA, Pfeiffer W, Schwartz T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Gateway Computing Environments Workshop (GCE): New Orleans, LA, USA, pp 1–8.
- Miyakawa K, Takama R, Kawabe I, Kariya S, Yamauchi T. (2010). A new automatic subsurface gas monitoring system for seismogeochimical studies, installed in Haruno Borehole, Shizuoka Prefecture, Central Japan. *Pure Appl Geophys* **167**: 1561–1575.
- Mohapatra BR, Dinardo O, Gould WD, Koren DW. (2010). Biochemical and genomic facets on the dissimilatory reduction of radionuclides by microorganisms—a review. *Miner Eng* **23**: 591–599.
- Nanjo I, Amano Y, Iwatsuki T, Kunimaru T, Murakami H, Hosoya S *et al.* (2012). Development of a Groundwater Monitoring System at Horonobe Underground Research Center. Available at: <http://jolissrch-inter.tokai-sc.jaea.go.jp/pdfdata/JAEA-Research-2011-048.pdf>.
- Nawrocki EP. (2009). *Structural RNA Homology Search and Alignment Using Covariance Models*. Washington University School of Medicine: St Louis, MO, USA.
- Nawrocki EP, Kolbe DL, Eddy SR. (2009). Infernal 1.0: inference of RNA alignments. *Bioinformatics* **25**: 1335–1337.

- Parkes RJ, Cragg B, Roussel E, Webster G, Weightman A, Sass H. (2014). A review of prokaryotic populations and processes in sub-seafloor sediments, including biosphere: geosphere interactions. *Mar Geol* **352**: 409–425.
- Pedersen K. (1997). Microbial life in granitic rock. *FEMS Microbiol Rev* **20**: 399–414.
- Pedersen K. (2000). Exploration of deep intraterrestrial microbial life: Current perspectives. *FEMS Microbiol Lett* **185**: 9–16.
- Pedersen K. (2005). Microorganisms and Their Influence on Radionuclide Migration in Igneous Rock Environments. *J Nucl Radiochem Sci* **6**: 11–15.
- Pedersen K. (2013). Metabolic activity of subterranean microbial communities in deep granitic groundwater supplemented with methane and H₂. *ISME J* **7**: 839–849.
- Peng Y, Leung HCM, Yiu SM, Chin FYL. (2012). IDBA-UD: a *de novo* assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**: 1420–1428.
- Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I *et al.* (2016). Genomic resolution of a cold subsurface aquifer community provides metabolic insights for novel microbes adapted to high CO₂ concentrations. *Environ Microbiol* **19**: 459–474.
- Probst AJ, Weinmaier T, Raymann K, Perras A, Emerson JB, Rattei T *et al.* (2014). Biology of a widespread uncultivated archaeon that contributes to carbon fixation in the subsurface. *Nat Commun* **5**: 1–13.
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J *et al.* (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* **35**: 7188–7196.
- Puigdomeneck I, Gurban I, Laaksoharju M, Luukkonen A, Löfman J, Pitkänen P *et al.* (2001). *Hydrochemical Stability of Groundwaters Surrounding a Spent Nuclear Fuel Repository in a 100 000 Year Perspective*. Helsinki, Finland. Available at: http://www.iaea.org/inis/collection/NCLCollectionStore/_Public/33/020/33020732.pdf?r=1.
- Rajala P, Raulio M, Sohlberg E, Bomberg M. (2015). Microbially induced corrosion of carbon steel in deep groundwater environment. *Front Microbiol* **6**: 1–13.
- Raveh-Sadka T, Thomas BC, Singh A, Firek B, Brooks B, Castelle CJ *et al.* (2015). Gut bacteria are rarely shared by co-hospitalized premature infants, regardless of necrotizing enterocolitis development. *Elife* **4**: e05477.
- Rudolph C, Wanner G, Huber R. (2001). Natural communities of novel archaea and bacteria growing in cold sulfurous springs with a string-of-pearls-like morphology. *Appl Environ Microbiol* **67**: 2336–2344.
- Sakurai K, Yoshikawa H. (2012). Isolation and identification of bacteria able to form biofilms from deep subsurface environments. *J Nucl Sci Technol* **49**: 287–292.
- Salinero K, Keller K, Feil WS, Feil H, Trong S, Di Bartolo G *et al.* (2009). Metabolic analysis of the soil microbe *Dechloromonas aromatica* str. RCB: indications of a surprisingly complex life-style and cryptic anaerobic pathways for aromatic degradation. *BMC Genomics* **10**: 351.
- Sasamoto H, Arthur RC, Hama K. (2011). Interpretation of undisturbed hydrogeochemical conditions in Neogene sediments of the Horonobe area, Hokkaido, Japan. *Appl Geochem* **26**: 1464–1477.
- Sasamoto H, Yamamoto N, Miyakawa K, Mizuno T. (2014). Data of groundwater chemistry obtained in the Horonobe Underground Research Laboratory Project (2011–2013). Available at: <http://jolissrch-inter.tokai-sc.jaea.go.jp/pdfdata/JAEA-Data-Code-2014-033.pdf>.
- Schattner P, Brooks AN, Lowe TM. (2005). The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res* **33**: W686–W689.
- Schut GJ, Adams MWW. (2009). The iron-hydrogenase of *Thermotoga maritima* utilizes ferredoxin and NADH synergistically: a new perspective on anaerobic hydrogen production. *J Bacteriol* **191**: 4451–4457.
- Sharon I, Morowitz MJ, Thomas BC, Costello EK, Relman DA, Banfield JF. (2013). Time series community genomics analysis reveals rapid shifts in bacterial species, strains, and phage during infant gut colonization. *Genome Res* **23**: 111–120.
- Shimizu S, Akiyama M, Ishijima Y, Hama K, Kunimaru T, Naganuma T. (2006). Molecular characterization of microbial communities in fault-bordered aquifers in the Miocene formation of northernmost Japan. *Geobiology* **4**: 203–213.
- Sieber JR, McInerney MJ, Gunsalus RP. (2012). Genomic insights into syntrophy: the paradigm for anaerobic metabolic cooperation. *Annu Rev Microbiol* **66**: 429–452.
- Stackebrandt E. (2014). The Family Acidimicrobiaceae. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson FL (eds). *The Prokaryotes*. Springer: Berlin, Heidelberg, Germany, pp 5–12.
- Stamatakis A. (2006). RAxML-VI-HPG: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.
- Suzek BE, Huang H, McCarvey P, Mazumder R, Wu CH. (2007). UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics* **23**: 1282–1288.
- Suzuki Y, Kelly SD, Kemner KM, Banfield JF. (2002). Nanometre-size products of uranium bioreduction. *Nature* **419**: 134.
- Suzuki Y, Mukai H, Ishimura T, Yokoyama TD, Sakata S, Hirata T *et al.* (2016). Formation and geological sequestration of uranium nanoparticles in deep granitic aquifer. *Sci Rep* **6**: 22701.
- Tachi Y, Yotsuji K, Seida Y, Yui M. (2011). Diffusion and sorption of Cs⁺, I[–] and HTO in samples of the argillaceous Wakkanai Formation from the Horonobe URL, Japan: Clay-based modeling approach. *Geochim Cosmochim Acta* **75**: 6742–6759.
- Tamamura S, Akatsuka M, Ikawa R, Koshigai M, Shimizu S, Ueno A *et al.* (2014). Origin of methane dissolved in formation waters in the Koetoi Formation through to the alluvium in northwestern part of Hokkaido, Japan. *Chikyukagaku* **48**: 39–50.
- Tatusova T, Ciufu S, Fedorov B, O'Neill K, Tolstoy I. (2014). RefSeq microbial genomes database: new representation and annotation strategy. *Nucleic Acids Res* **42**: D553–D559.
- Tremblay P, Zhang T, Dar SA, Leang C, Lovley DR. (2013). The Rnf complex of *Clostridium ljungdahlii* is a proton-translocating ferredoxin:NAD⁺ oxidoreductase essential for autotrophic growth. *MBio* **4**: e00406–e00412.
- Vinsot A, Appelo CAJ, Lundy M, Wechner S, Lettry Y, Lerouge C *et al.* (2014). *In situ* diffusion test of hydrogen gas in the Opalinus Clay. In: Geological Society, Vol 400. Special Publications: London, UK, pp 563–578.

- Wall JD, Krumholz LR. (2006). Uranium reduction. *Annu Rev Microbiol* **60**: 149–166.
- Wang X-J, Yang J, Chen X-P, Sun G-X, Zhu Y-G. (2009). Phylogenetic diversity of dissimilatory ferric iron reducers in paddy soil of Hunan, South China. *J Soils Sediments* **9**: 568–577.
- Weber KA, Achenbach LA, Coates JD. (2006). Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat Rev Microbiol* **4**: 752–764.
- Williams KH, Long PE, Davis JA, Wilkins MJ, N'Guessan AL, Steefel CI et al. (2011). Acetate availability and its influence on sustainable bioremediation of uranium-contaminated groundwater. *Geomicrobiol J* **28**: 519–539.
- Williamson AJ, Morris K, Shaw S, Byrne JM, Boothman C, Lloyd JR. (2013). Microbial reduction of Fe(III) under alkaline conditions relevant to geological disposal. *Appl Environ Microbiol* **79**: 3320–3326.
- Wolf PG, Biswas A, Morales SE, Greening C, Gaskins R. (2016). H₂ metabolism is widespread and diverse among human colonic microbes. *Gut Microbes* **7**: 235–245.
- Wood GE, Haydock AK, Leigh JA. (2003). Function and regulation of the formate dehydrogenase genes of the methanogenic archaeon *Methanococcus maripaludis*. *J Bacteriol* **185**: 2548–2554.
- Wouters K, Moors H, Boven P, Leys N. (2013). Evidence and characteristics of a diverse and metabolically active microbial community in deep subsurface clay borehole water. *FEMS Microbiol Ecol* **86**: 458–473.
- Wrighton KC, Castelle CJ, Varaljay VA, Satagopan S, Brown CT, Wilkins MJ et al. (2016). RubisCO of a nucleoside pathway known from Archaea is found in diverse uncultivated phyla in bacteria. *ISME J* **10**: 2702–2714.
- Wrighton KC, Thrash JC, Melnyk RA, Bigi JP, Byrne-Bailey KG, Remis JP et al. (2011). Evidence for direct electron transfer by a Gram-positive bacterium isolated from a microbial fuel cell. *Appl Environ Microbiol* **77**: 7633–7639.
- Wu WM, Carley J, Green SJ, Luo J, Kelly SD, Van Nostrand J et al. (2010). Effects of nitrate on the stability of uranium in a bioreduced region of the subsurface. *Environ Sci Technol* **44**: 5104–5111.
- Wu X, Holmfeldt K, Hubalek V, Lundin D, Åström M, Bertilsson S et al. (2015). Microbial metagenomes from three aquifers in the Fennoscandian shield terrestrial deep biosphere reveal metabolic partitioning among populations. *ISME J* **10**: 1192–1203.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)